

AD-A274 070

ATION PAGE

Form Approved

OMB No. 070-0188

2

average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. SUBJECT USE ONLY (Leave blank)		2. REPORT DATE 3 Dec 1993		3. REPORT TYPE AND DATES COVERED Final Report, May 91 - April 92	
4. TITLE AND SUBTITLE Antifouling Agents From Eelgrass: Identification, Characterization and Mode of Action				5. FUNDING NUMBERS N00014-91-J-1569	
6. AUTHOR(S) Richard C. Zimmerman and Hewson Swift					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Dept. Molecular Genetics & Cell Biology The University of Chicago Chicago IL 60637				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy St. Arlington VA 22217-5000				10. SPONSORING/MONITORING AGENCY REPORT NUMBER NTIS CRA&I <input checked="" type="checkbox"/> DTIC TAB <input type="checkbox"/> Unannounced <input type="checkbox"/> Justification	
11. SUPPLEMENTARY NOTES				By _____ Distribution /	
12a. DISTRIBUTION/AVAILABILITY STATEMENT Distribution Unlimited				12b. DISTRIBUTION CODES Dist Avail and/or Special A-1	
13. ABSTRACT (Maximum 200 words) This research has focused on the isolation and structural characterization of natural, non-toxic antifoulants produced by the eelgrass <u>Zostera marina</u> L. Chemical purification of a single active agent from fractions produced by aqueous-organic phase partitioning, size filtration chromatography and HPLC was directed by microbial attachment assays. Structural characterization of the isolated agent was accomplished by high resolution nuclear magnetic resonance and mass spectrometry. The isolated compound and several analogs were synthesized. Laboratory bioassays and short term (<10 day) field experiments indicate that the natural agent and at least of the one synthetic analogs show significant antifouling activity against marine bacteria, algal spores, benthic diatoms, tube worms and barnacles. Evaluation of their antifouling potential continues. A patent application describing the use of these compounds as non-toxic antifouling agents is in preparation.					
14. SUBJECT TERMS Biofouling, Natural Products, Eelgrass, Antifouling				15. NUMBER OF PAGES	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL		

93 12 21 134

93-30794

DTIC  
ELECTE  
DEC 23 1993

## Zimmerman & Swift, Antifouling Agents from Eelgrass

### GOALS:

The overall goals of this research have been to identify naturally-occurring, non-toxic antifoulant produced by the marine angiosperm, Zostera marina L. (eelgrass) and to characterize the mode of action of these agents in preventing attachment of fouling organisms.

### NEAR TERM OBJECTIVES:

Our objectives have focused on the isolation and structural characterization of the active agent(s), chemical synthesis of the isolated compound and several structural analogs, and evaluation of the antifouling effectiveness of the compounds. We have also screened extracts from other macrophyte taxa for the presence of antifouling activity and collected preliminary data on the antifouling effectiveness of the agent against hard foulers in the field.

### APPROACH:

Purification of a single active agent, using a sequence of aqueous-organic phase partitioning, size filtration chromatography and high pressure liquid chromatography (HPLC), was directed by a bacterial attachment bioassay. Structural characterization of the agent was accomplished by high resolution nuclear magnetic resonance and mass spectrometry.

## Zimmerman & Swift, Antifouling Agents from Eelgrass

### TASKS COMPLETED:

The active agent isolated from eelgrass has been purified and characterized structurally. This compound, along with several analogs, have been synthesized. Evaluation of the antifouling potential of these compounds continues. A patent application is in preparation for the use of the natural compound and synthetic analogs as non-polluting antifouling agents.

### RESULTS:

The synthetic compounds showed similar activity to the agent purified from eelgrass. Bacterial attachment to slides treated with one of the synthetic agents decreased exponentially with concentration (Fig. 1). The  $IC_{50}$  (concentration causing a 50% inhibition in bacterial density) was about  $1 \mu g \text{ ml}^{-1}$  ( $12 \mu g \text{ cm}^{-2}$  of treated surface). Despite the strong antifouling effect, there was no evidence of toxicity to bacterial growth in liquid culture or on agar at concentrations up to 10x higher than the  $IC_{50}$  dose. Thus, the therapeutic index ( $IC_{50}/LD_{50}$ ) remains undefined but extremely high ( $\geq 100$ ).

Soluble extracts were prepared from other local macrophytes and screened for antifouling activity. Of the 8 spp examined, including representative taxa from all major macrophyte groups, only extracts of Zostera marina showed significant activity (Fig. 2). Thus, the activity exhibited by the eelgrass extract was not a widespread phenomenon.

## Zimmerman & Swift, Antifouling Agents from Eelgrass

Preliminary field data indicate that the crude eelgrass extract effectively inhibited attachment of spirorbid worms (polychaetes) and colonial ascidians to unglazed ceramic tiles, while it was not effective against solitary tunicates (Fig. 3). No barnacles settled on any tiles during the course of this experiment.

The crude extract and a synthetic analog were also effective against barnacle larvae in laboratory attachment assays. The  $IC_{50}$  against barnacle attachment was about  $1 \mu\text{g/ml}$  ( $12 \mu\text{g/cm}^2$  of treated sfc.), similar to that demonstrated for bacteria (Fig. 4). As with bacteria, these agents were not toxic to barnacle larvae at concentrations that inhibited attachment.

### ACCOMPLISHMENTS:

The chemical agent isolated from eelgrass leaves has been purified and characterized structurally. It is effective at preventing attachment of a wide range of fouling organisms, from bacteria to barnacles, at concentrations that are orders of magnitude below the lethal dose. The agent is structurally simple, and has proven easy to synthesize from readily available and inexpensive substrates. Several manuscripts and a patent application are in preparation from this work.

**Zimmerman & Swift, Antifouling Agents from Eelgrass**

**PUBLICATIONS SUPPORTED BY THIS RESEARCH:**

**Kopczak, C.D., Zimmerman, R.C. and Kremer, J.N. 1991.**

**Variation in Nitrogen Physiology and Growth Among  
Geographically Isolated Populations of the Giant Kelp,  
Macrocystis pyrifera (Phaeophyta). J. Phycol. 27:148-158.**

**Zimmerman, R.C. and Alberte, R.S. 1992. Regulation of Carbon  
Partitioning in the Seagrass. Zostera marina L. (eelgrass).  
I. Assimilate Export Rates, Enzyme Activities and the Effect  
of Root Anoxia on Carbon Transport. Plant Physiol. In review.**

**Todd, J.T., Zimmerman, R.C. Dea, C.H., Crews, P., and Alberte,  
R.S. 1992. Isolation of natural antifouling agents from  
Zostera marina L. (eelgrass) and the antifouling activity of  
synthetic analogs. Phytochemistry. In prep.**

**Zimmerman, R.C. Todd, J., Dea, C.H. Baier, R., Crews, P., and  
Alberte, R.S. 1992. Mechanisms of Fouling Avoidance in  
Zostera marina L. (eelgrass). Mar. Biol. In prep.**

**PATENTS RESULTING FROM THIS RESEARCH:**

**Zimmerman, R.C. and Alberte, R.S. In Prep. Antifouling properties  
of natural zosteric acid isolated from eelgrass, and  
synthetic analogs. Arch Development Corp., Univ. Chicago,  
Chicago, IL 60637.**




Figure 1. Effect of concentration of a synthetic antifouling agent on attachment density of bacteria in laboratory assay. The  $IC_{50}$  concentration is about  $1 \mu g ml^{-1}$  ( $12 \mu g cm^{-2}$  of treated surface).




Figure 2. Results from bacterial attachment assays with extracts prepared for different marine macrophytes. Slides treated with solvent controls are indicated by open circles (O), slides treated with extract are indicated by filled circles (●).

Zimmerman & Swift, Antifouling Agents from Eelgrass




Figure 3. Effect of the crude eelgrass extract on settlement density of 3 invertebrate taxa in the field. Solvent control tiles are indicated by the open circles (O), extract-treated tiles are indicated by filled circles (●).






Figure 4. Effect of concentration of a synthetic antifouling agent on barnacle settlement in the laboratory. The  $\text{IC}_{50}$  for barnacles was about  $1 \mu\text{g ml}^{-1}$  ( $12 \mu\text{g cm}^{-2}$  of treated surface), similar to that for marine bacteria.

### **Distribution List for Final Reports**

**Attach a copy of the REPORT DOCUMENTATION PAGE (DD FORM 1473) to your final report as the first page and mail two copies (including the postcard labelled DTIC FORM 50 ) to:**

**Defense Technical Information Center  
Building 5, Cameron Station  
Alexandria, VA 22314**

**This allows other investigators to obtain copies of your report directly from DTIC. DTIC will fill out the postcard DTIC ACCESSION NOTICE (DTIC FORM 50) and return it to you with their number for your report. When you refer people to DTIC to get a copy of your report, give this number to expedite the request.**

**Mail one copy to each of the following and attach this very page to the back of your report - otherwise the folks below will think they have mistakenly received a copy meant for the Molecular Biology Program ):**

- |   |   |
|---|---|
| <b>(a) Dr. Michael Marron<br/>ONR Code 1141<br/>Molecular Biology Program<br/>800 N. Quincy Street<br/>Arlington, VA 22217-5000</b> | <b>(e) Director<br/>Chemical and Biological Sci Div<br/>Army Research Office<br/>P. O. Box 12211<br/>Research Triangle Park, NC 27709</b> |
| <b>(b) Administrative Grants Officer<br/>ONR Resident Representative<br/>(address varies - see copy of your<br/>grant/contract)</b> | <b>(f) Life Sciences Directorate<br/>Air Force Office of Scientific Res<br/>Bolling Air Force Base<br/>Washington, DC 20332</b>           |
| <b>(c) Director,<br/>Applied Research Directorate<br/>ONR Code 12<br/>800 N. Quincy Street<br/>Arlington, VA 22217-5000</b>         | <b>(g) Director<br/>Naval Research Laboratory<br/>Technical Information Div<br/>Code 2627<br/>Washington, DC 20375</b>                    |
| <b>(d) Director<br/>Office of Naval Technology<br/>Code 22<br/>800 N. Quincy Street<br/>Arlington, VA 22217-5000</b>                |   |